



Nano-mineralogy & -geochemistry of karst bauxites (Greece): Implications of the origin of the deposits

Gkamaletsos, Platon; Godelitsas, A.; Kasama, Takeshi; Göttlicher, J.; Steininger, R.

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Gkamaletsos, P., Godelitsas, A., Kasama, T., Göttlicher, J., & Steininger, R. (2016). *Nano-mineralogy & -geochemistry of karst bauxites (Greece): Implications of the origin of the deposits*. Abstract from V.M. Goldschmidt Conference 2016, Yokohama, Japan.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Nano-mineralogy & - geochemistry of karst bauxites (Greece): Implications of the origin of the deposits

P.N. GAMALETSOS^{1,2,3,*}, A. GODELITSAS², T.
KASAMA¹, J. GÖTTLICHER³, R. STEININGER³

¹ Center for Electron Nanoscopy, Technical
University of Denmark, Denmark
(*correspondence: plagka@dtu.dk)

² School of Science, University of Athens, Greece

³ ANKA Synchrotron Radiation Facility, Karlsruhe
Institute of Technology, Germany

Recent research with respect to the nano-mineralogy and -geochemistry of karst bauxites from Greece [1] is reviewed herein, while new insights into significant trace elements (including As, V, HFSE and REE) and the origin of these deposits are also presented [2].

The study of Fe-depleted (high grade) bauxite samples in microscale by SEM-EDS/WDS and Synchrotron Radiation (SR) measurements (μ -XRF & μ -XAFS) showed the presence of a Fe^{3+} - Cr^{3+} - AlOOH (diaspore), while the study in nanoscale by TEM-EDS & EELS revealed the existence of Fe^{3+} -oxide nanominerals and TiO_2 polymorph mineral nanoparticles [1]. It is argued that this Fe-Cr-diaspore (Fe^{3+} - Cr^{3+} - AlOOH) concerns a low-T authigenic phase, which was formed during diagenesis together with TiO_2 mineral micro- & nano-particles. The latter phases remain rather stable during epigenesis related to iron mobility and the formation of Fe nanominerals. The observed Fe nanominerals may be associated to epigenetic processes. It is also implied that younger karstification/weathering processes had minimal effect to Ti and Fe mineral nanoparticles and nanominerals [1]. New μ -XRF and μ -XAFS investigations of Fe-rich (low grade) samples revealed that As^{5+} is associated to Fe-Cr-Ti in pisoliths but, remarkably, not to S attributed to sulfides. Vanadium (most likely as V^{5+} , though contribution of V^{4+} cannot be excluded) is related to Ca, K, and possibly to Sc, whereas Ga is related to Al [2]. Detrital micro-sized chromite grains and positive anomalies of compatible elements (Ti, Cr, V, Ni, Sc) indicate a partial mafic-ultramafic origin (Tethys pre-Cretaceous ophiolites). The presence of HFS elements (Zr, Ce) and actinides (U, Th), whereas discrimination diagrams of contrasting elements (e.g., Th/Sc vs Zr/Sc) and geochemical fingerprints (including REE anomalies, i.e. Eu/Eu^* and Ce/Ce^*) strongly support a major contribution of acidic to intermediate igneous rocks derived, most likely, from pre-Jurassic peri-Tethyan large volcanic provinces.

[1] Gamaletsos P.N. et al., Am. Mineral. (under review).

[2] Gamaletsos P.N. et al., To be submitted to Ore Geol. Rev.